

## Quality and quantity: transitions in antimicrobial gland use for parasite defense

Article (Published Version)

Tranter, Christopher, Fernández-Marín, Hermógenes and Hughes, William O H (2015) Quality and quantity: transitions in antimicrobial gland use for parasite defense. *Ecology and Evolution*, 5 (24). pp. 5857-5868. ISSN 2045-7758

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/59999/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

## Quality and quantity: transitions in antimicrobial gland use for parasite defense

Christopher Tranter<sup>1</sup>, Hermógenes Fernández-Marín<sup>2</sup> & William O. H. Hughes<sup>1</sup>

<sup>1</sup>School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK

<sup>2</sup>Instituto de Investigaciones Científicas y Servicios de Alta Tecnología, Edificio 219, Panamá 5, Ciudad del Saber, Clayton, Panama City PO Box 0843-01105, Republic of Panama

### Keywords

Attini, disease resistance, fungus-growing ant, host–parasite evolution, leaf-cutting ant, metapleural gland.

### Correspondence

Christopher Tranter, School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK.  
Tel: 01273 678057;  
Fax: 01273 877586;  
E-mail: c.tranter@sussex.ac.uk

### Funding Information

Biotechnology and Biological Sciences Research Council (Grant/Award Number: 'BB/F016603/1').

Received: 23 September 2015; Revised: 9 October 2015; Accepted: 9 October 2015

**Ecology and Evolution 2015; 5(24): 5857–5868**

doi: 10.1002/ece3.1827

### Abstract

Parasites are a major force in evolution, and understanding how host life history affects parasite pressure and investment in disease resistance is a general problem in evolutionary biology. The threat of disease may be especially strong in social animals, and ants have evolved the unique metapleural gland (MG), which in many taxa produce antimicrobial compounds that have been argued to have been a key to their ecological success. However, the importance of the MG in the disease resistance of individual ants across ant taxa has not been examined directly. We investigate experimentally the importance of the MG for disease resistance in the fungus-growing ants, a group in which there is inter-specific variation in MG size and which has distinct transitions in life history. We find that more derived taxa rely more on the MG for disease resistance than more basal taxa and that there are a series of evolutionary transitions in the quality, quantity, and usage of the MG secretions, which correlate with transitions in life history. These shifts show how even small clades can exhibit substantial transitions in disease resistance investment, demonstrating that host–parasite relationships can be very dynamic and that targeted experimental, as well as large-scale, comparative studies can be valuable for identifying evolutionary transitions.

## Introduction

Parasites can inflict considerable costs on host organisms (Sheldon and Verhulst 1996; Rigby et al. 2002; Boots et al. 2004). This can result in a dramatic coevolutionary reshaping of the genotype, phenotype and overall life history of both hosts and parasites, and may have been a key influence in major evolutionary transitions including the evolution of sex and sociality (Hamilton et al. 1990; Møller et al. 1993; Boomsma et al. 2005). Understanding how the life history of hosts affects the selection strength on them from parasite pressure, and how this in turn leads to further changes in the evolutionary investment by hosts in costly disease resistance mechanisms, is an important problem in evolutionary biology (Schmid-Hempel 2001).

Group living is associated with a potential increase in parasite pressure, because a social lifestyle can facilitate the transmission of parasites within the group (Alexander

1974; Cote and Poulin 1995; Møller et al. 2001; Altizer et al. 2003). This is compounded in social insect colonies that have highly homeostatic nest environments and low levels of genetic variation within the colony (Schmid-Hempel 1994; Frank 1996; Calleri et al. 2006). However, in addition to individual-level immune defenses, social organisms are able to employ social defenses that have been termed “social immunity” in the broad sense (Dunbar 1991; Cremer et al. 2007; Wisenden et al. 2009; Otti et al. 2014). This can include behavioral defenses such as grooming, and the production and transfer of antimicrobial compounds (Rosengaus et al. 1998, 2004; Fernández-Marín et al. 2006; Yanagawa et al. 2008; Hamilton et al. 2011; Baracchi et al. 2012; Turnbull et al. 2012), which can mitigate or even outweigh the fitness cost from parasites for group-living animals (Rosengaus et al. 1998; Hughes et al. 2002; Ugelvig and Cremer 2007; Reber et al. 2011).

The threat of disease has led the most diverse group of social insects, the ants, to evolve a unique exocrine structure, the metapleural gland (MG), which varies in size between species and phenotypes, and in many taxa, produces an antimicrobial secretion that is spread over the cuticle either passively or, in some species, actively by grooming (Hölldobler and Wilson 1990; Bot and Boomsma 1996; Sumner et al. 2003; Fernández-Marín et al. 2006; Poulsen et al. 2006; Hughes et al. 2008, 2010; Yek and Mueller 2011). Consequently, ants are able to vary their level of investment in disease resistance both on an evolutionary timescale and as a short-term behavioral response to disease threat by active grooming of the secretion on to the cuticle. While the secretion from the metapleural gland can be antibacterial (Iizuka et al. 1979; Veal et al. 1992), effective at inhibiting fungal sporulation and growth, and helping ants resist parasites (Beattie et al. 1985, 1986; Bot et al. 2002; Graystock and Hughes 2011; Tranter et al. 2014), the gland is energetically costly to maintain (Poulsen et al. 2002b). Thus, its degree of use in different ant species can be used as a measure of their relative investment in disease resistance, and thus, to infer the strength of parasite pressure in different species with different life histories (Hughes et al. 2002; Poulsen et al. 2002a; Boomsma et al. 2005). While experimental blockage of the metapleural gland has been shown to increase the susceptibility of ants to fungal parasites (Poulsen et al. 2002b; Graystock and Hughes 2011; Tranter et al. 2014), the importance of the metapleural gland in disease resistance of individual ants from multiple different species has not previously been quantified directly.

One group of ants which has proved particularly powerful for comparative analyses of disease resistance are the fungus-growing ants (Attini) that form a monophyletic clade with well-developed MGs (Table S1; Mueller et al. 2001; Currie et al. 1999; Schultz and Brady 2008; Fernández-Marín et al. 2013). Fungus-growing ants have been most extensively investigated with respect to the defense of their fungal crop mutualist against a specialist parasitic fungus, *Escovopsis*. The ants achieve this using a combination of weeding, antimicrobial compounds from the MG, and antimicrobials produced by symbiotic actinomycete bacteria that the ants culture on their cuticles (Currie et al. 1999; Currie 2001a,b; Fernández-Marín et al. 2013). There appears to be a trade-off between these two sources of beneficial antimicrobials, with species that possess more of the *Escovopsis*-specific actinomycete defenses relying less heavily on the MG to protect their fungal crop (Fernández-Marín et al. 2013). However, while most attention has been focused on the mechanisms by which ants defend their fungal crop against parasites, the ants also need to defend themselves against disease. The antimicrobial compounds produced by the actinomycete

bacteria are thought to be specifically active against *Escovopsis* (Currie et al. 1999; 2003; Little et al., 2006; Pagnocca, 2012), and the ants rely for their own defense on mechanisms such as grooming and the MG. The antimicrobial activity of the MG secretion of leaf-cutting ants has been demonstrated in vitro (Bot et al. 2002), and the importance of the MG for defending both adults and brood against parasites has also been demonstrated in leaf-cutting ants experimentally (Poulsen et al. 2002a,b; Tranter et al. 2014; ). Previous comparative studies have shown that fungus-growing ants vary substantially in the size of their MG, the chemical composition of its secretion and their use of active MG grooming to spread the secretion, with the more evolutionary-derived *Atta* and *Acromyrmex* leaf-cutting ants that live in larger, more complex societies, having particularly large MG (Fernández-Marín et al. 2006, 2009, 2013; Hughes et al. 2008; Adams et al. 2012; Vieira et al. 2012b). However, whether these evolutionary transitions in MG size and secretion result in differences in the disease resistance of individual attine ants themselves has not previously directly quantified. In this study, we used MG use as a measure of investment into disease resistance within the evolutionary framework of the fungus-growing ants. Specifically, we test the hypotheses that when compared to more evolutionarily basal species, the more derived species with larger and more complex societies will show the following: (1) greater reliance on functioning MG glands for their own disease resistance; (2) greater active MG self-grooming rates in response to a parasite threat; and (3) more powerful antifungal components within the chemical makeup of MG secretions.

## Methods

### Study species

We studied six species of Neotropical attine ants spanning the major phylogenetic divisions within the clade: two species of leaf-cutting ant (*Atta colombica* and *Acromyrmex echinator*), three species of higher attines (*Sericomyrmex amabilis*, *Trachymyrmex cornetzi*, and *Trachymyrmex* sp10), and one species of lower attine (*Apterostigma pilosum*). These species represent a spectrum of both the agricultural systems employed within the fungus-growing ants (from the use of fresh leaves in *Acromyrmex* and *Atta*, to the use of other plant material in *Trachymyrmex* and *Sericomyrmex*, through to the use of general detritus or insect carcasses in *Apterostigma*), as well as the full range of colony sizes, social complexity, and phylogenetic ancestry. Colonies were collected in and around Gamboa, Panama in June 2013, and maintained at 80% relative humidity and 27°C on a 12-h light/dark cycle. Colonies were fed

twice per week with fresh privet (*Ligustrum* sp.) leaves for the two leaf-cutting species or chopped flower petals and oat flakes for the other four species, and provided with water ad libitum. As the experimental parasite, we used the entomopathogenic fungus *Metarhizium pingshaense* (KVL02-73; which was originally isolated from soil in Gamboa near a leaf-cutting ant nest; Hughes et al. 2004). *Metarhizium* is a ubiquitous, generalist fungal pathogen which parasitizes a wide range of insects, including attine and other ants, but which is unlikely because of its generalist nature to have coevolved to overcome the specific defenses of attine ants (Sanchez-Pena and Thorvilson 1992; Quiroz et al. 1996; Hughes et al. 2004; De Zarzuela et al. 2007; Castilho et al. 2010; Reber and Chapuisat 2011; Ribeiro et al. 2012).

### Experiment 1: the effect of gland blockage and fungal exposure on survival

Twenty worker ants were selected from just inside the nest entrance of each of six colonies of *S. amabilis*, *T. cornetzi*, *T. sp10*, and *A. pilosum*, 40 ants from six colonies of *A. echinator*, and 80 ants from three colonies of *A. colombica*. Trials involving *A. echinator* and *A. colombica* were conducted as two separate cohorts and data later pooled (see Fig. S1). Ants used within each species were a similar size, with medium cuticular coloration, and hence inferred age (Armitage and Boomsma 2010). For the polymorphic leaf-cutting ant species, we used workers of similar size to the other attines (0.9–1.4 mm head width). Half of the ants from each colony had their MG blocked using quick-drying nail varnish, while the other half received a control treatment of nail varnish applied to the pronotum. Nail varnish was checked daily and remained intact on all ants treated for the course of the experiment. After 24 h, each of these groups then had either a suspension of *Metarhizium* conidia in 0.05% Triton-X or a control solution of 0.05% Triton-X applied topically to the mesosoma with a micropipette. Treatment volumes were standardized for body size between species, and conidia concentrations were approximately the LD<sub>50</sub> for the species based on pilot studies ( $5 \times 10^6$  conidia per mL for leaf-cutting ant species and  $5 \times 10^5$  conidia per mL for the other species; Table S4). This design involved a total of 120 ants for each of *S. amabilis*, *T. cornetzi*, *T. sp10*, and *A. pilosum* (30 ants per species for each of the four treatment groups), and 240 ants for *A. colombica* and *A. echinator* (60 ants per species for each of the four treatment groups). After treatment, each ant was placed in a plastic pot (diameter: 35 mm, height: 70 mm) supplied with cotton balls soaked in 20% sucrose solution and water and kept at 70% relative humidity and 26°C. Ant mortality was recorded for 14 days. Cadavers were immediately removed and surface-

sterilized (Siegel 2012), and then kept in a Petri dish with moistened filter paper for an additional 14 days to allow the sporulation of fungi.

### Experiment 2: the effect of simulated fungal exposure on grooming rates

Twelve ants from each species (two individuals from each of six colonies for *A. echinator*, *S. amabilis*, *T. cornetzi*, *T. sp10*, *A. pilosum*, and four individuals from three colonies for *A. colombica*) were observed in a 40-mm Petri dish for 30 min with two nestmates. Incidences of allogrooming, self-grooming, and metapleural gland grooming of the focal ant were recorded (Altmann 1974). This process was repeated for 12 additional ants per species, but with each ant receiving a standardized treatment of dry, unscented, talcum powder (magnesium silicate) applied evenly to the dorsal mesosoma and gaster with a fine brush, to induce grooming, prior to observation for 30 min with two nestmates. We used talcum powder because some of the colonies were small and talcum powder particles, which are similar in size to fungal conidia and act as a nonpathogenic stimulant of antiparasite defensive behavior in ants without incurring the mortality that would result from application of fungal conidia (Fernández-Marín et al. 2006; Morelos-Juárez et al. 2010; Tranter et al. 2015).

### Experiment 3: chemical inhibition of fungal growth

Six isolated chemical compounds that have been previously identified by Vieira et al. (2012b) as major constituents of attine MG secretions were tested individually for their effect on the conidia viability of the entomopathogenic fungus *Metarhizium pingshaense*. The compounds tested were indole, skatole, methyl oleate (oleic acid), 2-nonanone, phenylacetic acid, and methyl-3-indoleacetate (indoleacetic acid; See Table S2 for percentage compositions found by Vieira et al. 2012b), as well as acetone solvent control, bleach (NaClO) positive control, and ddH<sub>2</sub>O negative control. Each individual compound was tested 10 times at five different concentrations based on the maximal amounts found in the MG of adult *Atta* workers (See Table S4 for summary of dilutions). A conidia solution of  $1 \times 10^5$  conidia per mL was prepared from freshly sporulating *M. pingshaense* plates. 450 agar plates were prepared with selective media (Sabouraud dextrose agar [SDA] with 0.05 g/L streptomycin sulfate and 0.1 g/L chloramphenicol antibiotics, and 0.1 g/L dodin which inhibits the growth of other fungi but not *Metarhizium*; Shah et al. 2005) in 50-mm Petri dishes and stored sealed at 4°C until use. 500 µL of the *Metarhizium* conidia

solution ( $5 \times 10^6$  conidia per mL) was applied evenly over the surface of the Petri dish and left for 10 min to allow excess liquid to dry. A single, 6-mm diameter piece of sterile plastic tubing was placed carefully onto the center of the surface of the agar plates, and 20  $\mu$ L of test solution was applied in the center of this with the plastic cylinder acting as a well to restrict distribution to a defined area. The cylinder was left in place for 5 min to allow the compound to infuse the media, before the location of the treated area was marked on the underside of the Petri dish and the cylinder removed. The Petri dish was then sealed with parafilm and placed in an incubator at 32°C overnight. 12 h later, the percentage of conidia producing a germ tube longer than the conidia diameter (Siegel 2012; conidia viability) was counted for a standardized area (complete area visible in the microscope eyepiece at 400 $\times$  magnification) within the section where the compound was applied and also an untreated area outside the marked test area, equidistant with the edge of the Petri dish. A further 60 h later, the plates were photographed from above and the average radius of any zone of inhibition, as characterized by an area around the marked test section free from fungal growth, was recorded (Fig. S2).

### Statistical analysis

The effects of *Metarhizium* exposure, gland closure, and ant species, on ant survival in Experiment 1 were analyzed using Cox proportional hazards regression models. Colony-of-origin and cohort for the leaf-cutting ant trials were included in the models to account for the structured nature of the data, but were not statistically significant ( $P > 0.05$  in all cases). Pairwise comparisons were made with Kaplan–Meier tests using the Breslow statistic. The numbers of cadavers sporulating with *Metarhizium* for ants with blocked or unblocked glands were examined for each species with Chi-squared tests. Grooming rates following exposure to talcum powder in Experiment 2 were analyzed using a general linear mixed model with a gamma distribution and log-link function; colony-of-origin was included as a random factor but was not statistically significant ( $P > 0.05$  in all cases). Nonsignificant interaction terms were removed stepwise to obtain the minimum adequate model in each case. Pairwise comparisons were conducted between treatments within each species, and between species for each treatment. The effects of compound and dose in Experiment 3 on the size of fungal inhibition zones were analyzed using a generalized linear model with a gamma distribution and log-link function. Multiple comparisons were controlled for in all analyses using the sequential Bonferroni adjustment. All analyses were performed in IBM SPSS v21 Armonk, NY: IBM Corp.

## Results


### Experiment 1: the effect of gland blockage and fungal exposure on survival

Overall, there were significant effects of species and interaction between blockage and fungal treatments on survival ( $Wald = 17$ ,  $df = 5$ ,  $P = 0.005$ ;  $Wald = 9.98$ ,  $df = 1$ ,  $P = 0.02$ , respectively). Ants from all six species showed a significant reduction in their survival when treated with the *Metarhizium* parasite (Fig. 1; Table S3). *A. colombica*, *A. echinator*, and *S. amabilis* all showed a significant reduction in resistance to the parasite when their MG was blocked, while the resistance of ants with blocked and unblocked MG was nearly identical in both *T. cornetzi* and *A. pilosum*, and there was also no significant effect of MG blockage on the resistance of *T. sp10* (Fig. 1; Table S3). There was no significant effect of colony on survival in any of the species ( $P > 0.05$ ; Table S3). None of the control ants sporulated with *Metarhizium*. Of those *Metarhizium*-exposed ants that died, significantly more of the cadavers sporulated with *Metarhizium* when they had blocked glands compared to those where the MG was functional ( $\chi^2 = 47.8$ ,  $df = 5$ ,  $P < 0.001$ ). This difference was present in *A. colombica*, *A. echinator*, and *S. amabilis*, but not in *T. cornetzi*, *T. sp10*, or *A. pilosum* (Fig. 1).


### Experiment 2: the effect of simulated fungal exposure on grooming rates

Contact rates between ants differed between species ( $F_{1,137} = 20.8$ ,  $P < 0.001$ ), reflecting interspecific differences in the general activity levels of the ants, but there was no effect of talcum powder application or interaction between effects on activity ( $F_{1,132} = 0.42$ ,  $P = 0.52$ ;  $F_{5,132} = 1.12$ ,  $P = 0.36$ ; Fig. 2A). There was, however, a significant interaction between the effects of ant species and talcum powder treatment on MG grooming ( $F_{5,132} = 9.67$ ,  $P = 0.014$ ). *A. colombica* exhibited the highest rate of MG grooming by far, with *A. echinator* and *S. amabilis* also conducting higher levels of MG grooming compared to the *Trachymyrmex* and *Apterostigma* species which exhibited little or no MG grooming (Fig. 2B). In spite of the relatively small sample sizes for some species, there was overall also a significant interaction between species and talcum powder treatment on the rates of self-grooming ( $F_{5,132} = 3$ ,  $P = 0.014$ ), with *T. sp10* and *A. pilosum* both self-grooming significantly more when talcum powder was applied to them and baseline levels of self-grooming being highest in *T. cornetzi* (Fig. 2C). Only *S. amabilis* allogroomed significantly more when exposed to talcum powder treatment ( $F_{1,132} = 5.5$ ,  $P = 0.02$ ; Fig. 2D), and there was no overall effect of species or





**Figure 1.** Survival over 2 weeks of (A) *Apterostigma pilosum*, (B) *Trachymyrmex cornetzi*, (C) *Trachymyrmex sp10*, (D) *Sericomyrmex amabilis*, (E) *Acromyrmex echinator*, and (F) *Atta colombica* attine ants after treatment with either *Metarhizium pingshaense* fungal parasite (solid line) or control solution (dashed line), and with their antimicrobial-producing metapleural glands either experimentally blocked (closed circles) or functional (open circles). Different letters beside lines indicate treatments which differed significantly from one another within species at  $P < 0.05$ . Inset graphs show the proportions of cadavers of *Metarhizium*-exposed ants that sporulated with *Metarhizium* for ants of each species with either their glands blocked ("B": dark bars) or functional ("F": light bars). Species in which the frequency of sporulation differed significantly between ants with blocked and functional glands at  $P < 0.05$  are marked with an asterisk (\*).



**Figure 2.** The mean  $\pm$  SE frequencies of (A) contact, (B) metapleural gland grooming, (C) self-grooming, and (D) allogrooming in 30 min for six species of attine ants (*Atta colombica*, *Acromyrmex echinator*, *Sericomyrmex amabilis*, *Trachymyrmex sp10*, *Trachymyrmex cornetzi*, and *Apterostigma pilosum*) either treated with talcum powder (dark bars) or untreated (light bars). Significant differences between treated and untreated ants at  $P < 0.05$  for each species are indicated with an asterisk. Species which differed significantly from one another at  $P < 0.05$  are indicated by different letters, a, b, c for untreated ants, or x, y, z for treatment ants.

interaction between factors for allogrooming ( $F_{5,132} = 0.4$ ,  $P = 0.86$ ;  $F_{5,132} = 0.79$ ,  $P = 0.56$ ).

### Experiment 3: chemical inhibition of fungal growth


There was a significant interaction between the compound tested and the dose applied on the size of the zone in which *Metarhizium* fungal growth was inhibited ( $\chi^2 = 194.5$ ,  $df = 32$ ,  $P < 0.001$ ), and on the number of fungal conidia that were viable ( $\chi^2 = 575.4$ ,  $df = 32$ ,  $P < 0.001$ ). Phenylacetic acid consistently produced the largest reductions in spore viability, especially at higher doses (Fig. 3A). It also produced the largest inhibition zone in the highest dose, where it was generally comparable to, or even more effective than, bleach in its antifungal activity (Fig. 3B).

### Discussion

Exposure to the fungal parasite *Metarhizium pingshaense* significantly reduced the survival of the six species of ants

in this study, and there were species-specific differences between the resistance of ants when their MGs were blocked, compared with when they were left functional. There were also differences in MG grooming rates and the antifungal activity of chemical components of the MG secretions between species.

The leaf-cutting ants *A. colombica* and *A. echinator*, as well as the higher attine *S. amabilis*, which also has relatively large colony sizes, all demonstrated a consistently greater reliance on MG use for protection against fungal pathogens compared with the more basal *Trachymyrmex* and *Apterostigma* species (Fig. 4). The resistance of leaf-cutting ants and *Sericomyrmex* to the parasite was significantly reduced when their MG glands were blocked. Additionally, of those ants that died, significantly more of the cadavers sporulated with the parasite when the MG was blocked. This highlights the importance of the gland in sterilizing the ant's cuticle and promoting their survival through fungistatic or fungicidal effects. The leaf-cutting ants and *S. amabilis* all also showed higher rates of MG grooming behavior, and their secretions contained chemicals with stronger antifungal activity, including more




**Figure 3.** The mean  $\pm$  SE reduction in viability of conidia of the *Metarhizium pingshaense* fungal parasite (A) and size of growth inhibition zone produced (B), for six chemicals from the metapleural gland secretion (phenylacetic acid, 2-nonanone, indole, skatole, methyl-3-indolacetate, methyl oleate), and bleach positive control, acetone, and hexane solvent control, with each compound applied at five concentrations (1 – lowest dose – 5 highest: see Table S4 for details). A ddH<sub>2</sub>O negative control produced negligible effect and was omitted from the graph. For doses 1, 3, and 5 where pairwise comparisons were performed, different letters indicate chemicals which differed significantly from one another at  $P < 0.05$ .

acidic compounds phenylacetic acid and methyl oleate (Do Nascimento and Schoeters 1996; Yek et al. 2012). This was particularly evident in *Atta colombica*, which was the only species in which the highly antifungal

phenylacetic acid has been identified (Vieira et al. 2012b; Fernández-Marín et al. 2015). Additionally, the strength of this effect, and reliance on the MG, seems to be compounded by the efficacy of some of the MG secretions





**Figure 4.** Phylogenetic patterns of metapleural gland usage in the attine ants. Graphs show for the six species studied here (highlighted in dark gray boxes): (1) the hazard ratio from blockage of the antibiotic-producing metapleural gland for ants exposed to the *Metarhizium* fungal parasite; (2) the mean  $\pm$  SE frequency of active grooming of the metapleural gland in a 30-min period; (3) the effectiveness of the overall metapleural gland secretion at reducing *Metarhizium* conidia viability, based on the sum of the average effectiveness of each chemical weighted by its representation in the secretion (Tables S4, S5); (4) the size of the metapleural gland reservoir (bulla width) relative to body size (pronotum width; data from Hughes et al. (2008) and includes data for *Apterostigma collare*, not *Apterostigma pilosum* as in our study). The number of ant symbols shown next to species name represents the typical number of workers per colony in orders of magnitude.

themselves; not only do the leaf-cutting ants and *S. amabilis* use their glands more actively, but also the compounds in their secretions are more effective antifungal agents. These findings support our predictions that the more derived leaf-cutting ant species and *S. amabilis* with their larger and more complex societies are considerably more reliant on their MG compared to the more basal species.

The importance of the MG in resisting disease varied across the attine phylogeny providing evidence for the existence of a series of four evolutionary transitions in MG use (Fig. 4). First, between *Apterostigma* and *Trachymyrmex* sp10, we see a small increase in the use of the MG and consequent increased importance of the MG in resisting disease. Second, there is then a small increase in MG size and antifungal activity, and a large increase in MG grooming and the importance of the MG for disease resistance, between *Trachymyrmex* and *Sericomyrmex*. Third, there is a substantial increase in MG gland size between *Sericomyrmex* and the leaf-cutting ant species. Fourth, there is a transition between *Acromyrmex* and *Atta*, with *Atta* producing a more powerful suite of antifungal chemicals within their MG secretions, which they produce more of and groom much more actively than *Acromyrmex* (Fig. 4). There is no evidence of a trade-off between different aspects of individual self-directed MG use, but rather a general trend across the phylogeny toward the MG becoming more effective and important

in disease resistance. Although there is a significant phylogenetic distance between *Apterostigma* and *Trachymyrmex*, the differences in MG use between these species were less dramatic than the transitions observed between species higher up the phylogeny, in which there are also more dramatic changes in colony size and complexity. Further studies on more basal species intermediate between *Apterostigma* and *Trachymyrmex* are needed to confirm this and allow formal quantitative phylogenetically controlled analyses, as well as to reveal whether the MG in these species may instead have a role in defense against different microbes.

These differences in MG use, in combination with transitions in other morphological and behavioral attributes such as gardening of the fungal crop or hitchhiking on harvested leaf material, may reflect changes in host–parasite interactions within the attine clade, notably, as a result of specialization of the fungal mutualism and the appearance of polymorphic workers (Hughes et al. 2008; Schultz and Brady 2008; Fernández-Marín et al. 2009; Griffiths and Hughes 2010; Vieira et al. 2012a). Within the Attini, the leaf-cutting species, and *Atta* especially, are the most morphologically and behaviorally specialized taxa (Hart et al. 2002; Evison and Ratnieks 2007), and, as we demonstrate here, are especially well adapted for parasite defense. The presence of discreet castes allows for a differentiation in MG morphology in the higher attine species, with the smallest castes having relatively larger

MG reservoirs (Hughes *et al.* 2008). This may aid in parasite resistance through allowing investment into MG defense in those individuals that may benefit from them the most. In leaf-cutting ants these are the minors that tend to the fungus, clean leaf fragments, and brood (Currie 2001a,b; Hughes *et al.* 2008; Griffiths and Hughes 2010). It may be that these derived characters have allowed for greater investment into costly disease resistance through improved resource acquisition; for example, the fungal gardens cultivated on fresh vegetation by *Atta* and *Acromyrmex* ants may provide a more nutritious and reliable food source.

Additionally, MG use within the group maps broadly with colony size between the species (Fig. 4; Table S1). Larger colonies have a greater workforce, greater resource acquisition, and are a more robust entity that can provide more stable nest conditions (Kunz 1982; Rosengren *et al.* 1987; Jeanne and Nordheim 1996; Anderson and Ratnieks 1999; Jones and Oldroyd 2006; Jeanson *et al.* 2007). However, larger group size may also result in increases in parasite pressure through the same stable and favorable nest conditions and increased colony longevity, which promote parasite transmission (Schmid-Hempel 1998; Poulin 2007) and may “sample” more of the environment which increases the chance of contracting parasites (Wilson 1971; Rosengren *et al.* 1987; Sherman *et al.* 1988; Hölldobler and Wilson 1990; Tschinkel 1991; Schmid-Hempel 1998; Zahn 1999; Poulin 2007). An alternative explanation is therefore that the larger colonies of leaf-cutting ants may be exposed to greater parasite pressures than lower attine species and thus invest more in disease resistance in order to mitigate this increased cost (Hart and Ratnieks 1998; Hart and Ratnieks 2001; Hughes *et al.* 2002; Naug and Camazine 2002; Poulsen *et al.* 2002b; Fernández-Marín *et al.* 2006, 2009; Tranter *et al.* 2014).

Our results show how changes in the antifungal activity and use of MG secretions affect the resistance of individual ants to parasites, and demonstrate that even relatively small clades can exhibit substantial transitions in investment into disease resistance mechanisms. This highlights how dynamic the evolutionary relationships between host and parasite can be, and demonstrates the value of targeted experimental studies on multiple species for identifying and understanding evolutionary transitions in host–parasite relationships.

## Acknowledgments

We are grateful to Allen Herre and the Smithsonian Tropical Research Institute for facilities in Gamboa, the Autoridad Nacional del Ambiente (ANAM) for permission to collect and export colonies, and the BBSRC for funding.

## Conflict of Interest

This work was supported by the Biotechnology and Biological Sciences Research Council. The authors of this paper declare no conflict of interests. All applicable international, national, and institutional guidelines for the care and use of animals were followed. This article does not contain any studies with human participants performed by any of the authors.

## References

- Adams, R. M. M., T. H. Jones, A. W. Jeter, H. H. De Fine Licht, T. R. Schultz, and D. R. Nash. 2012. A comparative study of exocrine gland chemistry in *Trachymyrmex* and *Sericomyrmex* fungus-growing ants. *Biochem. Syst. Ecol.* 40:91–97.
- Alexander, R. D. R. 1974. The evolution of social behavior. *Annu. Rev. Ecol. Evol. Syst.* 5:325–383.
- Altizer, S., C. Nunn, and P. Thrall. 2003. Social organization and parasite risk in mammals: integrating theory and empirical studies. *Annu. Rev. Ecol. Evol. Syst.* 34:517–547.
- Altmann, J. 1974. Observational study of behavior: sampling methods. *Behaviour* 49:227–267.
- Anderson, C., and F. Ratnieks. 1999. Task partitioning in insect societies. I. Effect of colony size on queueing delay and colony ergonomic efficiency. *Am. Nat.* 154:521–535.
- Armitage, S. A. O., and J. J. Boomsma. 2010. The effects of age and social interactions on innate immunity in a leaf-cutting ant. *J. Insect Physiol.* 56:780–787.
- Baracchi, D., G. Mazza, and S. Turillazzi. 2012. From individual to collective immunity: the role of the venom as antimicrobial agent in the Stenogastrinae wasp societies. *J. Insect Physiol.* 58:188–193.
- Beattie, A. J., C. Turnbull, T. Hough, S. Jobson, and R. B. Knox. 1985. The vulnerability of pollen and fungal spores to ant secretions: evidence and some evolutionary implications. *Am. J. Bot.* 1:606–614.
- Beattie, A. J., C. Turnbull, T. Hough, and R. B. Knox. 1986. Antibiotic production: a possible function for the metapleural glands of ants (Hymenoptera: Formicidae). *Ann. Entomol. Soc. Am.* 79:448–450.
- Boomsma, J. J., P. Schmid-Hempel, and W. O. H. Hughes. 2005. Life histories and parasite pressure across the major groups of social insects. Pp. 139–176 in M. Fellowes, G. Holloway and J. Rolff, eds. *Insect evolutionary ecology*. Wallingford, UK, CABI Publishing.
- Boots, M., P. J. Hudson, and A. Sasaki. 2004. Large shifts in pathogen virulence relate to host population structure. *Science* 303:842–844.
- Bot, A. N. M., and J. J. Boomsma. 1996. Variable metapleural gland size-allometries in *Acromyrmex* leafcutter ants (Hymenoptera: Formicidae). *J. Kansas Entomol. Soc.* 69:375–383.

- Bot, A. N. M., D. Ortius-Lechner, K. Finster, R. Maile, and J. J. Boomsma. 2002. Variable sensitivity of fungi and bacteria to compounds produced by the metapleural glands of leaf-cutting ants. *Insectes Soc.* 49:363–370.
- Calleri, D. V., E. McGrail Reid, R. B. Rosengaus, E. L. Vargo, and J. F. A. Traniello. 2006. Inbreeding and disease resistance in a social insect: effects of heterozygosity on immunocompetence in the termite *Zootermopsis angusticollis*. *Proc. R. Soc. Lond. B* 273:2633–2640.
- Castilho, A. M. C., M. E. Fraga, L. de, E. Aguiar-menezes, R. da, C. A. Rosa, et al. 2010. Seleção de isolados de *Metarhizium anisopliae* e *Beauveria bassiana* patogênicos a soldados de *Atta bisphaerica* e *Atta sexdens rubropilosa* em condições de laboratório. *Ciência Rural* 40:1243–1249.
- Cote, I. M., and R. Poulin. 1995. Parasitism and group size in social animals: a meta-analysis. *Behav. Ecol.* 6:159–165.
- Cremer, S., S. A. O. Armitage, and P. Schmid-Hempel. 2007. Social immunity. *Curr. Biol.* 17:693–702.
- Currie, C. R. 2001a. A community of ants, fungi, and bacteria: a multilateral approach to studying symbiosis. *Annu. Rev. Microbiol.* 55:357–380.
- Currie, C. R. 2001b. Weeding and grooming of pathogens in agriculture by ants. *Proc. R. Soc. Lond. B* 268:1033–1039.
- Currie, C. R., J. Scott, R. Summerbell, and D. Malloch. 1999. Fungus-growing ants use antibiotic-producing bacteria to control garden parasites. *Nature* 398:701–704.
- Currie, C. R., B. Wong, A. E. Stuart, T. R. Schultz, S. Rehner, U. G. Mueller, et al. 2003. Ancient tripartite coevolution in the attine ant-microbe symbiosis. *Science* 299:386–388.
- De Zarzuela, M. F. M., A. E. de Carvalho Campos-Farinha, P. C. Kruppa, and E. Gonzalez. 2007. Evaluation of urban ants (Hymenoptera: Formicidae) as vectors of microorganisms in residential and industrial environments: II Fungi. *Sociobiology* 50:653–658.
- Do Nascimento, R., and E. Schoeters. 1996. Chemistry of metapleural gland secretions of three attine ants, *Atta sexdens rubropilosa*, *Atta cephalotes*, and *Acromyrmex octospinosus* (Hymenoptera: Formicidae). *J. Chem. Ecol.* 22:987–1000.
- Dunbar, R. I. M. 1991. Functional significance of social grooming in primates. *Folia Primatol.* 57:121–131.
- Evison, S. E. F., and F. L. W. Ratnieks. 2007. New role for majors in *Atta leafcutter* ants. *Ecol. Entomol.* 32:451–454.
- Fernández-Marín, H., J. K. Zimmerman, S. A. Rehner, and W. T. Wcislo. 2006. Active use of the metapleural glands by ants in controlling fungal infection. *Proc. R. Soc. Lond. B* 273:1689–1695.
- Fernández-Marín, H., J. K. Zimmerman, D. R. Nash, J. J. Boomsma, and W. T. Wcislo. 2009. Reduced biological control and enhanced chemical pest management in the evolution of fungus farming in ants. *Proc. R. Soc. Lond. B* 276:2263–2269.
- Fernández-Marín, H., G. Bruner, E. B. Gomez, D. R. Nash, J. J. Boomsma, and W. T. Wcislo. 2013. Dynamic disease management in *Trachymyrmex* fungus-growing ants (Attini: Formicidae). *Am. Nat.* 181:571–582.
- Fernández-Marín, H., D. R. Nash, S. Higginbotham, C. Estrada, J. van Zweeken, P. d'Ettorre, et al. 2015. Functional role of phenylacetic acid from metapleural gland secretions in controlling fungal pathogens in evolutionarily derived leaf-cutting ants. *Proc. R. Soc. Lond. B* 282:20150212.
- Frank, S. A. 1996. Models of parasite virulence. *Q. Rev. Biol.* 71:37–78.
- Graystock, P., and W. O. H. Hughes. 2011. Disease resistance in a weaver ant, *Polyrhachis dives*, and the role of antibiotic-producing glands. *Behav. Ecol. Sociobiol.* 65:2319–2327.
- Griffiths, H. M., and W. O. H. Hughes. 2010. Hitchhiking and the removal of microbial contaminants by the leaf-cutting ant *Atta colombica*. *Ecol. Entomol.* 35:529–537.
- Hamilton, W. D., R. Axelrod, and R. Tanese. 1990. Sexual reproduction as an adaptation to resist parasites (a review). *Proc. Natl Acad. Sci. USA* 87:3566–3573.
- Hamilton, C., B. T. Lejeune, and R. B. Rosengaus. 2011. Trophallaxis and prophylaxis: social immunity in the carpenter ant *Camponotus pennsylvanicus*. *Biol. Lett.* 7:89–92.
- Hart, A. G., and F. L. W. Ratnieks. 1998. Waste management in the leaf-cutting ant *Atta colombica*. *Behav. Ecol.* 13:224–231.
- Hart, A. G., and F. L. W. Ratnieks. 2001. Task partitioning, division of labour and nest compartmentalisation collectively isolate hazardous waste in the leafcutting ant *Atta cephalotes*. *Behav. Ecol. Sociobiol.* 49:387–392.
- Hart, A. G., C. Anderson, and F. L. W. Ratnieks. 2002. Task partitioning in leafcutting ants. *Acta Ethol.* 5:1–11.
- Hölldobler, B., and E. O. Wilson. 1990. The ants. Belknap Press, Cambridge, MA.
- Hughes, W. O. H., J. Eilenberg, and J. J. Boomsma. 2002. Trade-offs in group living: transmission and disease resistance in leaf-cutting ants. *Proc. R. Soc. Lond. B* 269:1811–1819.
- Hughes, W. O. H., L. Thomsen, J. Eilenberg, and J. J. Boomsma. 2004. Diversity of entomopathogenic fungi near leaf-cutting ant nests in a neotropical forest, with particular reference to *Metarhizium anisopliae* var. *anisopliae*. *J. Invertebr. Pathol.* 85:46–53.
- Hughes, W. O. H., R. Pagliarini, H. B. Madsen, M. B. Dijkstra, and J. J. Boomsma. 2008. Antimicrobial defense shows an abrupt evolutionary transition in the fungus-growing ants. *Evolution* 62:1252–1257.
- Hughes, W. O. H., A. N. M. Bot, and J. J. Boomsma. 2010. Caste-specific expression of genetic variation in the size of antibiotic-producing glands of leaf-cutting ants. *Proc. R. Soc. Lond. B* 277:609–615.
- Iizuka, T., T. Iwadare, and K. Orito. 1979. Antibacterial activity of myrmicacin and related compounds on pathogenic bacteria in silkworm larvae, *Streptococcus faecalis* AD-4. *J. Fac. Agric. Hokkaido Univ.* 59:262–266.

- Jeanne, R. L., and E. V. Nordheim. 1996. Productivity in a social wasp: per capita output increases with swarm size. *Behav. Ecol.* 7:43–48.
- Jeanson, R., J. H. Fewell, R. Gorelick, and S. M. Bertram. 2007. Emergence of increased division of labor as a function of group size. *Behav. Ecol. Sociobiol.* 62:289–298.
- Jones, J. C., and B. P. Oldroyd. 2006. Nest thermoregulation in social insects. *Adv. Insect Phys.* 33:153–191.
- Kunz, T. H. 1982. *Ecology of bats*. Springer, US, Boston, MA.
- Little, A. E., T. Murakami, U. G. Mueller, and C. R., Currie. 2006. Defending against parasites: fungus-growing ants combine specialized behaviours and microbial symbionts to protect their fungus gardens. *Biol. Lett.* 2:12–16.
- Møller, A., D. Reija, K. Allander, R. Dufva, and K. Allander. 1993. Parasites and the evolution of host social behavior. Pp. 65–102 in C. T. Snowdon and M. Milinski, eds. *Advances in the study of behavior*, Vol. 22. Academic Press, Waltham, USA.
- Møller, A., S. Merino, C. Brown, and R. Robertson. 2001. Immune defense and host sociality: a comparative study of swallows and martins. *Am. Nat.* 158:136–145.
- Morelos-Juárez, C., T. N. Walker, J. F. S. Lopes, and W. O. H. Hughes. 2010. Ant farmers practice proactive personal hygiene to protect their fungus crop. *Curr. Biol.* 20:553–554.
- Mueller, U. G., T. R. Schultz, C. R. Currie, R. M. M. Adams, and D. Malloch. 2001. The origin of the attine ant-fungus mutualism. *Q. Rev. Biol.* 76:169–197.
- Naug, D., and S. Camazine. 2002. The role of colony organization on pathogen transmission in social insects. *J. Theor. Biol.* 215:427–439.
- Ortius-Lechner, D., R. Maile, E. D. Morgan, and J. J. Boomsma. 2000. Metapleural gland secretion of the leaf-cutter ant *Acromyrmex octospinosus*: new compounds and their functional significance. *J. Chem. Ecol.* 26:1667–1683.
- Otti, O., S. Tragust, and H. Feldhaar. 2014. Unifying external and internal immune defences. *Trends Ecol. Evol.* 29:625–634.
- Pagnocca, F. 2012. Specialized fungal parasites and opportunistic fungi in gardens of attine ants. *Psyche*. 1–9.
- Poulin, R. 2007. *Evolutionary ecology of parasites*. Princeton Univ. Press, Princeton, NJ.
- Poulsen, M., A. N. M. Bot, C. R. Currie, and J. J. Boomsma. 2002a. Mutualistic bacteria and a possible trade-off between alternative defence mechanisms in *Acromyrmex* leaf-cutting ants. *Insectes Soc.* 49:15–19.
- Poulsen, M., A. N. M. Bot, M. G. Nielsen, and J. J. Boomsma. 2002b. Experimental evidence for the costs and hygienic significance of the antibiotic metapleural gland secretion in leaf-cutting ants. *Behav. Ecol. Sociobiol.* 52:151–157.
- Poulsen, M., W. O. H. Hughes, and J. J. Boomsma. 2006. Differential resistance and the importance of antibiotic production in *Acromyrmex echinator* leaf-cutting ant castes towards the entomopathogenic fungus *Aspergillus nomius*. *Insectes Soc.* 53:349–355.
- Quiroz, L., G. Carrion, and J. Valenzuela. 1996. Hongos entomopatogenos de las hormigas arrieras *Atta mexicana* en México. *Rev. Mex. Micol.* 12:41–48.
- Reber, A., and M. Chapuisat. 2011. Diversity, prevalence and virulence of fungal entomopathogens in colonies of the ant *Formica selysi*. *Insectes Soc.* 2:231–239.
- Reber, A., J. Purcell, S. D. Buechel, P. Buri, and M. Chapuisat. 2011. The expression and impact of antifungal grooming in ants. *J. Evol. Biol.* 24:954–964.
- Ribeiro, M. M. R., K. D. Amaral, V. E. Seide, B. M. R. Souza, T. M. C. Della Lucia, M. C. M. Kasuya, et al. 2012. Diversity of fungi associated with *Atta bisphaerica* (Hymenoptera : Formicidae): the activity of *Aspergillus ochraceus* and *Beauveria bassiana*. *Psyche* 2012:1–6.
- Rigby, M. C., R. F. Hechinger, and L. Stevens. 2002. Why should parasite resistance be costly? *Trends Parasitol.* 18:116–120.
- Rosengaus, R. B., A. B. Maxmen, L. E. Coates, and J. F. A. Traniello. 1998. Disease resistance: a benefit of sociality in the dampwood termite *Zootermopsis angusticollis* (Isoptera: Termopsidae). *Behav. Ecol. Sociobiol.* 44:125–134.
- Rosengaus, R. B., J. F. A. Traniello, M. L. Lefebvre, and A. B. Maxmen. 2004. Fungistatic activity of the sternal gland secretion of the dampwood termite *Zootermopsis angusticollis*. *Insectes Soc.* 51:259–264.
- Rosengren, R., W. Fortelius, K. Lindström, and A. Luther. 1987. Phenology and causation of nest heating and thermoregulation in red wood ants of the *Formica rufa* group studied in coniferous forest habitats in southern Finland. *Ann. Zool. Fenn.* 24:147–155.
- Sanchez-Pena, S. R., and H. G. Thorvilson. 1992. Two fungi infecting red imported fire ant founding queens from Texas. *Southwest. Entomol.* 17:181–182.
- Schmid-Hempel, P. 1994. Infection and colony variability in social insects. *Philos. Trans. R. Soc. Lond. B.* 346:313–321.
- Schmid-Hempel, P. 1998. *Parasites in social insects*. Princeton Univ. Press, Princeton, NJ.
- Schmid-Hempel, P. 2011. *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford Univ. Press, Oxford, U.K.
- Schultz, T. R., and S. G. Brady. 2008. Major evolutionary transitions in ant agriculture. *Proc. Natl Acad. Sci. USA* 105:5435–5440.
- Shah, F. A., C. S. Wang, and T. M. Butt. 2005. Nutrition influences growth and virulence of the insect-pathogenic fungus *Metarhizium anisopliae*. *FEMS Microbiol. Lett.* 251:259–266.
- Sheldon, B., and S. Verhulst. 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* 11:317–321.
- Sherman, P. W. W., T. D. D. Seeley, and H. K. K. Reeve. 1988. Parasites, pathogens, and polyandry in social Hymenoptera. *Am. Nat.* 131:602–610.



- Siegel, J. P. 2012. Manual of techniques in invertebrate pathology. Elsevier, Amsterdam.
- Sumner, S., W. O. H. Hughes, and J. J. Boomsma. 2003. Evidence for differential selection and potential adaptive evolution in the worker caste of an inquiline social parasite. *Behav. Ecol. Sociobiol.* 54:256–263.
- Tranter, C., P. Graystock, C. Shaw, J. F. S. Lopes, and W. O. H. Hughes. 2014. Sanitizing the fortress: protection of ant brood and nest material by worker antibiotics. *Behav. Ecol. Sociobiol.* 68:499–507.
- Tranter, C., L. LeFevre, S. E. F. Evison, and W. O. H. Hughes. 2015. Threat detection: contextual recognition and response to parasites by ants. *Behav. Ecol.* 26:396–405.
- Tschinkel, W. R. 1991. Insect sociometry, a field in search of data. *Insectes Soc.* 38:77–82.
- Turnbull, C., H. Caravan, T. Chapman, D. Nipperess, S. Dennison, M. Schwarz, et al. 2012. Antifungal activity in thrips soldiers suggests a dual role for this caste. *Biol. Lett.* 8:526–529.
- Ugelvig, L. V., and S. Cremer. 2007. Social prophylaxis: group interaction promotes collective immunity in ant colonies. *Curr. Biol.* 17:1967–1971.
- Veal, D. A., J. E. Trimble, and A. J. Beattie. 1992. Antimicrobial properties of secretions from the metapleural glands of *Myrmecia gulosa* (the Australian bull ant). *J. Appl. Microbiol.* 72:188–194.
- Vieira, A. S., O. C. Bueno, and M. I. Camargo-Mathias. 2012a. Morphophysiological differences between the metapleural glands of fungus-growing and non-fungus-growing ants (Hymenoptera, Formicidae). *PLoS ONE* 7:e43570.
- Vieira, A. S., E. D. Morgan, F. P. Drijfhout, and M. I. Camargo-Mathias. 2012b. Chemical composition of metapleural gland secretions of fungus-growing and non-fungus-growing ants. *J. Chem. Ecol.* 10:1289–1297.
- Wilson, E. O. 1971. The insect societies. Harvard Univ. Press, Cambridge, MA.
- Wisenden, B., C. Goater, and C. James, 2009. Behavioral defenses against parasites and pathogens. Pp. 151–168 in B.G. Kapoor, Z. Giacomo, C. Perriere and A. Mathis, eds. Fish defenses: pathogens, parasites, predators. Science Publishers, Enfield, NH, USA.
- Yanagawa, A., F. Yokohari, and S. Shimizu. 2008. Defense mechanism of the termite, *Coptotermes formosanus* Shiraki, to entomopathogenic fungi. *J. Invertebr. Pathol.* 97:165–170.
- Yek, S. H., and U. G. Mueller. 2011. The metapleural gland of ants. *Biol. Rev.* 86:774–791.
- Yek, S. H., D. R. Nash, A. B. Jensen, and J. J. Boomsma. 2012. Regulation and specificity of antifungal metapleural gland secretion in leaf-cutting ants. *Proc. R. Soc. Lond. B* 279:4215–4222.
- Zahn, A. 1999. Reproductive success, colony size and roost temperature in attic-dwelling bat *Myotis myotis*. *J. Zool.* 247:275–280.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Figure S1.** Experimental groups and species cohort information for Exp 1.

**Figure S2.** Photos (top) of two plates (left: phenylacetic acid, right: bleach) showing zones around central application point in which growth of the *Metarhizium* fungal parasite was inhibited.

**Table S1.** Fungus growing ant life-history traits.

**Table S2.** MG compounds tested as a percentage of total secretion volume for Attine genera.

**Table S3.** Statistical results of survival analysis in six attine species with blocked or functional metapleural glands, and treated with *Metarhizium pinshaense* fungal parasite or control solution.

**Table S4.** Isolated MG secretions listed by their abundances as reported in GC-MS fractions (Do Nascimento and Schoeters 1996; Ortius-Lechner et al. 2000; Vieira et al. 2012b) and natural gland secretions from *Acromyrmex* ants.

**Table S5.** The natural abundances, individual antimicrobial activities, and calculated percentage antimicrobial activity for MG secretion compounds tested in Attine ants.

**Table S6.** Experiment 1: Survival of ants treated with Triton-X control or *Metarhizium* fungus with glands blocked or unblocked.